

## References

1. Wilczek HE, Jaremko G, Tyden G, Groth CG. Evolution of diabetic nephropathy in kidney grafts. Evidence that a simultaneously transplanted pancreas exerts a protective effect. *Transplantation* 1995; 59: 51–7.
2. Brayman KL, Stephanian E, Matas AJ, et al. Analysis of infectious complications occurring after solid-organ transplantation. *Arch Surg* 1992; 127: 38–47; discussion 47–8.
3. Kentouni-Noly JC, Cloix P, Martin X, Rabodonirina M, Lefrancois N, Sepetjan M. Analysis of nosocomial infections in renal transplant and pancreas transplant recipients. *Transplant Proc* 1994; 26: 284.
4. Pouteil-Noble C, Ecochard R, Landrison G, et al. Cytomegalovirus infection—an etiological factor for rejection? A prospective study in 242 renal transplant patients. *Transplantation* 1993; 55: 851–7.
5. Sollinger HW, Ploeg RJ, Eckhoff DE, et al. Two hundred consecutive simultaneous pancreas–kidney transplants with bladder drainage. *Surgery* 1993; 114: 736–43; discussion 743–4.
6. Am DA, Sollinger HW, Stratta RJ, Kalayoglu M, Pirsch JD, Belzer FO. Comparison between duodenal button and duodenal segment in pancreas transplantation. *Transplantation* 1989; 47: 120–2.
7. Nghiem DD, Corry RJ. Technique of simultaneous renal pancreatoduodenal transplantation with urinary drainage of pancreatic secretion. *Am J Surg* 1987; 153: 405–6.
8. Starzl TE, Fung J, Jordan M, et al. Kidney transplantation under FK 506. *JAMA* 1990; 264: 63–7.
9. Baldi A, Mourad M, Malaise J, Squifflet JP. Low urinary tract infections in simultaneous pancreas and kidney transplantation: comparison between bladder and enteric drainage of exocrine secretions. *Transplant Proc* 1995; 27: 3116–17.
10. Douzjian V, Abecassis MM, Cooper JL, Smith JL, Corry RJ. Incidence, management and significance of surgical complications after pancreatic transplantation. *Surg Gynecol Obstet* 1993; 177: 451–6.
11. Secchi A, Caldara R, La Rocca E, et al. Simultaneous kidney and pancreas transplantation at the San Raffaele Scientific Institute: clinical experience and results. *Clin Transplant* 1994; 8: 255–63.
12. Schulak JA, Mayes JT, Hricik DE. Combined kidney and pancreas transplantation. A safe and effective treatment for diabetic nephropathy. *Arch Surg* 1990; 125: 881–5.
13. Whelchel JD, Larse C, Olsen R, et al. Simultaneous pancreas kidney transplantation: initial experience of the Emory University transplant service. *Int Surg* 1994; 79: 98–102.
14. Lumberras C, Fernandez I, Velosa J, Munn S, Sterioff S, Paya CV. Infectious complications following pancreatic transplantation: incidence, microbiological and clinical characteristics, and outcome. *Clin Infect Dis* 1995; 20: 514–20.
15. Troppmann C, Gruessner AC, Benedetti E, Papalois BE, Sutherland DE, Gruessner RW. Positive duodenal segment cultures are not associated with increased surgical complications after whole organ, bladder–drained pancreas transplantation in three recipient categories. *Transplant Proc* 1995; 27: 3101–3.
16. Stratta RJ, Taylor RJ, Sindhi R, Sudan D, Castaldo P. Infection prophylaxis in pancreas transplant recipients. *Transplant Proc* 1995; 27: 3018–19.
17. Van der Pijl JW, Smets YF, de la Fuente R, Ringers J, Lemkes HH, van der Woude FJ. Urologic infections and problems after 50 simultaneous pancreas–kidney transplantations. *Transplant Proc* 1995; 27: 3105.
18. Benedetti E, Gruessner AC, Troppmann C, et al. Intra-abdominal fungal infections after pancreatic transplantation: incidence, treatment, and outcome. *J Am Coll Surg* 1996; 183: 307–16.
19. Abendroth D, La Greca G, Capalbo M, Illner WD, Landgraf R, Land W. Infections in simultaneous pancreas transplantation duct: occlusion vs bladder drainage. *Transplant Proc* 1992; 24: 927–8.
20. Sollinger HW, Messing EM, Eckhoff DE, et al. Urological complications in 210 consecutive simultaneous pancreas–kidney transplants with bladder drainage. *Ann Surg* 1993; 218: 561–8; discussion 568–70.
21. Ploeg RJ, Eckhoff DE, Am DA, et al. Urological complications and enteric conversion after pancreas transplantation with bladder drainage. *Transplant Proc* 1994; 26: 458–9.
22. Kuo PC, Johnson LB, Schweitzer EJ, Bartlett ST. Simultaneous pancreas/kidney transplantation—a comparison of enteric and bladder drainage of exocrine pancreatic secretions. *Transplantation* 1997; 63: 238–43.
23. Douzjian V, Rajagopalan PR. Primary enteric drainage of the pancreas allograft revisited. *J Am Coll Surg* 1997; 185: 471–5.
24. Ricordi C, Alejandro R, Angelico MC, et al. Human islet allografts in patients with type 2 diabetes undergoing liver transplantation. *Transplantation* 1997; 63: 473–5.

## In vitro activity of linezolid and eperezolid against anaerobic bacteria

*Clin Microbiol Infect* 1999; 5: 51–53

Charlotta Edlund<sup>1,2\*</sup>, Herin Oh<sup>1</sup> and Carl Erik Nord<sup>1</sup>

<sup>1</sup>Department of Immunology, Microbiology, Pathology and Infectious Diseases, Huddinge University Hospital, S-141 86 Huddinge; <sup>2</sup>University College of South Stockholm, Stockholm, Sweden

\*Tel: +46 8 58581139 Fax: +46 8 7113918 E-mail: charlotta.edlund@impi.ki.se

Accepted 22 June 1998

Linezolid (PNU-100766) and eperezolid (PNU-100592) belong to a new class of synthetic antimicrobial agents called the oxazolidinones [1]. These agents are highly active against many clinically important Gram-positive bacteria including methicillin-resistant staphylococci, vancomycin-resistant enterococci and penicillin-resistant pneumococci with MIC<sub>90</sub> values usually ranging from 0.5 to 4 mg/L [2,3]. Gram-negative aerobic microorganisms are generally not inhibited by these compounds. The oxazolidinones exhibit a unique mechanism of bacterial protein synthesis inhibition, resulting in mainly bacteriostatic activity [1].

Anaerobic bacteria are a common cause of serious infections. The anaerobic species which predominate in clinical infections include the *Bacteroides fragilis* group, *Clostridium* spp., *Fusobacterium nucleatum* and *Peptostreptococcus* spp. Treatment of anaerobic infections is often difficult, since many anaerobes are intrinsically resistant to certain antimicrobial agents such as aminoglycosides, trimethoprim, sulfamethoxazole, most quinolones and monobactams [4]. Resistance against tetracyclines is widespread, and many clinically important anaerobes exhibit resistance against clindamycin and several  $\beta$ -lactam agents. Resistance against cefoxitin, imipenem and metronidazole has been reported, although it is uncommon. Drugs commonly used in the treatment of anaerobic infections are  $\beta$ -lactam compounds, clindamycin, metronidazole and chloramphenicol [5].

The aim of the present study was to determine the in vitro activity of linezolid and eperezolid against 360 anaerobic bacterial strains isolated from human infections.

Three hundred and sixty anaerobic clinical isolates, including *Peptostreptococcus* spp. (50 strains), *Propionibacterium acnes* (30 strains), *Clostridium perfringens* (30 strains), *C. difficile* (50 strains), *B. fragilis* (50 strains), *Bacteroides*, *Porphyromonas*, and *Prevotella* spp. (100 strains) and fusobacteria (50 strains), collected at the Huddinge University Hospital, Stockholm, Sweden were tested. All strains were identified by biochemical tests and gas-liquid chromatographic analyses of metabolic end products according to the techniques described by Summanen et al. [6]. The strains were cultured in prereduced yeast glucose medium or in prereduced chopped meat broth with glucose. Linezolid and eperezolid were obtained from Pharmacia & Upjohn Inc., Kalamazoo, MI, USA. Fresh dilutions of each compound were prepared daily. The antimicrobial agents were suspended and diluted according to the manufacturers' instructions. Antimicrobial susceptibility tests were performed by the agar dilution method using

PDM-ASM agar (Biodisk, Stockholm, Sweden), with the addition of 5% defibrinated horse blood. Antimicrobial concentrations from 0.008 to 128 mg/L were obtained by incorporation of each substance when preparing the agar plates. The inocula consisted of 48-h broth cultures diluted in prereduced yeast glucose medium to a final inoculum of 10<sup>8</sup> CFU/mL. Then, 1.0–2.0  $\mu$ L was applied to the agar plates with a modified Steers replicator, resulting in approximately 10<sup>3</sup> CFU per spot. An agar plate without antimicrobial agent was always included as growth control. Agar plates were incubated in anaerobic jars (GasPak, BBL, Microbiology Systems, Cockeysville, MD, USA) for 48 h at 37°C. The MIC was defined as the lowest concentration that inhibited growth of organisms on the agar plates. A barely visible haze or the appearance of a single colony was disregarded. Three control strains were used to monitor the antimicrobial susceptibility test: *B. fragilis* ATCC 25285, *B. thetaiotaomicron* ATCC 29741 and *C. perfringens* ATCC 13124.

Ninety per cent of all tested *Peptostreptococcus* spp. (50 strains), *Propionibacterium acnes* (30 strains), *C. perfringens* (50 strains) and *C. difficile* (50 strains) were inhibited by  $\leq 2$  mg/L linezolid and eperezolid.

**Table 1** In vitro activity of linezolid and eperezolid against 360 anaerobic bacterial strains

Microorganism (no. of isolates tested) and antimicrobial agent	MIC (mg/L)		
	Range	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Peptostreptococcus</i> (50)			
Linezolid	0.25–2.0	1.0	2.0
Eperezolid	0.25–1.0	0.5	1.0
<i>Propionibacterium acnes</i> (30)			
Linezolid	0.25–1.0	0.5	0.5
Eperezolid	0.5–1.0	0.5	1.0
<i>Clostridium perfringens</i> (50)			
Linezolid	1.0–4.0	2.0	2.0
Eperezolid	1.0–2.0	2.0	2.0
<i>Clostridium difficile</i> (50)			
Linezolid	1.0–2.0	1.0	2.0
Eperezolid	0.5–1.0	1.0	1.0
<i>Bacteroides fragilis</i> (100)			
Linezolid	2.0–4.0	4.0	4.0
Eperezolid	2.0–32	8.0	16.0
<i>Bacteroides</i> , <i>Porphyromonas</i> , <i>Prevotella</i> (50)			
Linezolid	0.25–8.0	2.0	4.0
Eperezolid	0.125–8.0	2.0	8.0
<i>Fusobacterium</i> (30)			
Linezolid	0.25–8.0	0.5	8.0
Eperezolid	0.125–2.0	0.25	2.0

Linezolid showed higher activity (MIC<sub>90</sub> 4.0 mg/L) against *B. fragilis* (100 strains) compared to eperezolid (MIC<sub>90</sub> 16 mg/L). The MIC values (range, MIC<sub>50</sub> and MIC<sub>90</sub>) for the 360 tested strains of linezolid and eperezolid are shown in Table 1. Strains with intrinsic resistance to conventional anti-anaerobe agents, including three *C. difficile* strains and one *B. fragilis* strain resistant to clindamycin, seven *B. fragilis* strains resistant to cefoxitin, and one *B. splanchnicus* resistant to metronidazole, did not show any cross-resistance to the oxazolidinones.

The results of the present study are in accordance with previous reports on in vitro susceptibility to linezolid and eperezolid of anaerobic microorganisms [3]. Oral administration of linezolid (400 mg PO) and eperezolid (1000 mg PO) to healthy volunteers has earlier been reported to yield peak serum concentrations of 12.38 mg/L and 6.28 mg/L, respectively, while the trough concentrations were estimated to be 7.9 mg/L and 1.62 mg/L, respectively [7]. These serum levels of linezolid are well in excess of the MICs for anaerobic Gram-positive bacteria as well as for many anaerobic Gram-negative microorganisms. Concerning eperezolid, anaerobic Gram-positive bacteria are mainly susceptible to achieved serum levels. Linezolid has been reported to be active versus experimental *B. fragilis* soft tissue infections in mice [8]. Future clinical studies will show the potential of the oxazolidinones in the treatment of anaerobic infections. In conclusion, the new oxazolidinones linezolid and eperezolid have excellent activity against Gram-positive anaerobic

microorganisms, with linezolid also exhibiting activity against *Bacteroides* spp. These agents may be useful in the treatment and prophylaxis of anaerobic infections.

#### Acknowledgment

This work was supported by Pharmacia & Upjohn Inc., Kalamazoo, MI, USA.

#### References

1. Ford CW, Hamel JC, Stapert D, et al. Oxazolidinones: new antibacterial agents. *Trends Microbiol* 1996; 5: 196–200.
2. Jones RN, Johnson DM, Erwin ME. In vitro antimicrobial activities and spectra of U-100592 and U-100766, two novel fluorinated oxazolidinones. *Antimicrob Agents Chemother* 1996; 40: 720–6.
3. Zurenco GE, Yagi BH, Schaadt RD, et al. In vitro activities of U-100592 and U-100766, novel oxazolidinone antibacterial agents. *Antimicrob Agents Chemother* 1996; 40: 839–45.
4. Finegold S. Anaerobic infections in humans: an overview. *Anaerobe* 1995; 1: 3–9.
5. Hedberg M, Nord CE. Antimicrobial-resistant anaerobic bacteria in human infections. *Med Microbiol Lett* 1996; 5: 295–304.
6. Summanen P, Baron E, Citron D, Strong C, Wexler H, Finegold S. *Wadsworth anaerobic bacteriology manual*, 5th edn. Los Angeles, CA: Veterans Administration Wadsworth Medical Center, 1993.
7. Schaadt RD, Batts DH, Daley-Yates PT, Pawsey SD, Stalker DJ, Zurenco GE. Serum inhibitory titers and serum bactericidal titers for human subjects receiving multiple doses of the antibacterial oxazolidinones eperezolid and linezolid. *Diagn Microbiol Infect Dis* 1997; 28: 201–4.
8. Ford CW, Hamel JC, Wilsom DM, et al. The in vivo activities of U-100592 and U-100766, novel oxazolidinone antimicrobial agents, against experimental bacterial infections. *Antimicrob Agents Chemother* 1996; 40: 1508–13.

## Identification of methicillin-resistant *Staphylococcus aureus* by latex agglutination kits: performance with epidemic strains (EMRSA) and strains causing problems with latex agglutination methods

*Clin Microbiol Infect* 1999; 5: 53–56

Colin W. B. Walker\* and Derek F. J. Brown

Clinical Microbiology and Public Health Laboratory, Box 236, Addenbrooke's Hospital, Cambridge CB2 2QW, UK

\*Tel: +44 1223 257035 Fax: +44 1223 242775 E-mail:

Accepted 22 June 1998

Agglutination kits for the identification of *Staphylococcus aureus* have been available for a number of years as an alternative to the coagulase test. The tube coagulase

test, which detects the production of free coagulase, was considered to be the standard (sensitivity 95–98%) but it takes 4–24 h for completion [1,2]. A more simple